

Hindawi Publishing Corporation  
Journal of Inequalities and Applications  
Volume 2007, Article ID 87414, 8 pages  
doi:10.1155/2007/87414

## Research Article

# Subsequential Convergence Conditions

İbrahim Çanak, Ümit Totur, and Mehmet Dik

Received 27 April 2007; Accepted 19 August 2007

Recommended by Martin J. Bohner

Let  $(u_n)$  be a sequence of real numbers and let  $L$  be any  $(C, 1)$  regular limitable method. We prove that, under some assumptions, if a sequence  $(u_n)$  or its generator sequence  $(V_n^{(0)}(\Delta u))$  generated regularly by a sequence in a class  $\mathcal{A}$  of sequences is a subsequential convergence condition for  $L$ , then for any integer  $m \geq 1$ , the  $m$ th repeated arithmetic means of  $(V_n^{(0)}(\Delta u))$ ,  $(V_n^{(m)}(\Delta u))$ , generated regularly by a sequence in the class  $\mathcal{A}^{(m)}$ , is also a subsequential convergence condition for  $L$ .

Copyright © 2007 İbrahim Çanak et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## 1. Introduction

Let  $(u_n)$  be a sequence of real numbers. Let  $c_0$ ,  $\ell_\infty$ ,  $\mathcal{S}$ , and  $\mathcal{M}$  denote the space of sequences converging to 0, bounded, slowly oscillating, and moderately oscillating, respectively.

The classical control modulo of the oscillatory behavior of  $(u_n)$  is denoted by  $\omega_n^{(0)}(u) = n\Delta u_n$ , where  $\Delta u_n = u_n - u_{n-1}$  and  $u_{-1} = 0$  and the general control modulo of the oscillatory behavior of integer order  $m$  of  $(u_n)$  is defined [1] inductively by  $\omega_n^{(m)}(u) = \omega_n^{(m-1)}(u) - \sigma_n^{(1)}(\omega^{(m-1)}(u))$ , where  $\sigma_n^{(1)}(u) = (1/(n+1)) \sum_{k=0}^n u_k$ .

The Kronecker identity  $u_n - \sigma_n^{(1)}(u) = V_n^{(0)}(\Delta u)$ , where  $V_n^{(0)}(\Delta u) = (1/(n+1)) \sum_{k=0}^n k\Delta u_k$ , is well known and used in various steps of proofs of theorems. For each integer  $m \geq 1$  and for all nonnegative integers  $n$ , we inductively define sequences related to  $(u_n)$  such as  $V_n^{(m)}(\Delta u) = \sigma_n^{(1)}(V^{(m-1)}(\Delta u))$  and  $\sigma_n^{(m)}(u) = \sigma_n^{(1)}(\sigma^{(m-1)}(u))$ , where  $\sigma_n^{(0)}(u) = u_n$ .

Throughout this work, a different definition of slow oscillation better tailored for our purposes will be used. A sequence  $u = (u_n)$  is slowly oscillating [2] if  $\lim_{\lambda \rightarrow +\infty} \overline{\lim}_n \max_{n+1 \leq k \leq [\lambda n]} |u_k - u_n| = 0$ , where  $[\lambda n]$  denotes the integer part of  $\lambda n$ . See [3, 4] for more on slow oscillation. A sequence  $u = (u_n) \in \mathcal{S}$  if and only if  $(V_n^{(0)}(\Delta u)) \in \mathcal{S}$

and  $(V_n^{(0)}(\Delta u)) \in \ell_\infty$  (see [5]). A sequence  $u = (u_n)$  is moderately oscillating [2] if for  $\lambda > 1$ ,  $\overline{\lim}_n \max_{n+1 \leq k \leq [\lambda n]} |u_k - u_n| < \infty$ . It is proved in [5] that if a sequence  $u = (u_n) \in \mathcal{M}$ , then  $(V_n^{(0)}(\Delta u)) \in \ell_\infty$ .

A sequence  $u = (u_n)$  is Abel limitable to  $s$  if the limit  $\lim_{x \rightarrow 1^-} (1-x) \sum_{n=0}^{\infty} u_n x^n = s$  and  $(C, 1)$  limitable to  $s$  if  $\lim_n \sigma_n^{(1)}(u) = s$ .

Let  $L$  be any limitation method. If  $u = (u_n)$  is  $L$  limitable to  $s$ , we write  $L - \lim_n u_n = s$ . The limitation method  $L$  is said to be regular if  $\lim_n u_n = s$  implies  $L - \lim_n u_n = s$ . The limitation method  $L$  is said to be  $(C, 1)$  regular if  $L - \lim_n u_n = s$  implies  $L - \lim_n \sigma_n^{(1)}(u) = s$ . A sequence  $u = (u_n)$  is called subsequentially convergent [6] if there exists a finite interval  $I(u)$  such that all accumulation points of  $u = (u_n)$  are in  $I(u)$  and every point of  $I(u)$  is an accumulation point of  $u = (u_n)$ .

Let  $\mathcal{L}$  be any linear space of sequences and let  $\mathcal{A}$  be a subclass of  $\mathcal{L}$ . For each integer  $m \geq 1$ , define the class  $\mathcal{A}^{(m)} = \{(a_n^{(m)}) \mid a_n^{(m)} = \sum_{k=1}^n (a_k^{(m-1)}/k)\}$ , where  $(a_n^{(0)}) := (a_n) \in \mathcal{A}$ . Let  $u = (u_n) \in \mathcal{L}$ . If

$$u_n = a_n^{(m)} + \sum_{k=1}^n \frac{a_k^{(m)}}{k} \quad (1.1)$$

for some  $a^{(m)} = (a_n^{(m)}) \in \mathcal{A}^{(m)}$ , we say that the sequence  $(u_n)$  is regularly generated by the sequence  $(a_n^{(m)})$  and  $(a_n^{(m)})$  is called a generator of  $(u_n)$ . The class of all sequences regularly generated by sequences in  $\mathcal{A}^{(m)}$  is denoted by  $U(\mathcal{A}^{(m)})$ . We note that  $\mathcal{A}^{(0)} = \mathcal{A}$ .

Tauber [7] proved that an Abel limitable sequence  $u = (u_n)$  is convergent if

$$(\omega_n^{(0)}(u)) \in c_o. \quad (1.2)$$

A condition such as (1.2) is called a Tauberian condition, after A. Tauber.

Tauber [7] further proved that the condition

$$(\sigma_n^{(1)}(\omega^{(0)}(u))) \in c_o \quad (1.3)$$

is also a Tauberian condition. It was later shown by Littlewood [8] that the condition (1.2) could be replaced by

$$(\omega_n^{(0)}(u)) \in \ell_\infty. \quad (1.4)$$

Rényi [9] observed that the condition

$$(\sigma_n^{(1)}(\omega^{(0)}(u))) \in \ell_\infty \quad (1.5)$$

is no longer a Tauberian condition for Abel limitable method.

Stanojević [1] investigated behaviors of some subsequences of an Abel limitable sequence  $u = (u_n)$  adding a mild condition on  $(u_n)$ , together with (1.5).

Dik [6] obtained the following theorem.

THEOREM 1.1. *Let  $(u_n)$  be Abel limitable and  $\Delta V_n^{(0)}(\Delta u) = o(1)$ . If*

$$(V_n^{(0)}(\Delta u)) \in U(\mathcal{M}), \quad (1.6)$$

*then  $(u_n)$  is subsequentially convergent.*

Later several improvements of Dik's theorem were obtained.

A condition that subsequential convergence of  $(u_n)$  is recovered out of its Abel limitability is called a subsequential convergence condition.

We list the subsequential convergence conditions for Abel limitable method that (1.6) can be replaced by

- (i)  $(V_n^{(m)}(\Delta u)) \in U(\mathcal{M}^{(m)})$  (see [10]),
- (ii)  $(V_n^{(0)}(\Delta u)) \in U(\ell_\infty)$  (see [6]),
- (iii)  $(V_n^{(m)}(\Delta u)) \in U(\ell_\infty^{(m)})$  (see [10]),
- (iv)  $(u_n) \in U(\mathcal{M})$  (see [11]),
- (v)  $(u_n) \in U(\ell_\infty)$  (see [6]).

In this work, we prove that under the assumptions if a sequence  $(u_n)$  or its generator sequence  $(V_n^{(0)}(\Delta u))$  generated regularly by a sequence in a class  $\mathcal{A}$  of sequences is a subsequential convergence condition for a  $(C, 1)$  regular limitable method  $L$ , then for any integer  $m \geq 1$ , the  $m$ th repeated arithmetic means of  $(V_n^{(0)}(\Delta u))$ ,  $(V_n^{(m)}(\Delta u))$ , generated regularly by a sequence in the class  $\mathcal{A}^{(m)}$  is also a subsequential convergence condition for  $L$ .

## 2. Results

Throughout this section, we require  $L$  to be  $(C, 1)$  regular.

We prove the following theorems.

THEOREM 2.1. *For a sequence  $u = (u_n)$ , let  $L - \lim_n u_n = s$  and  $\Delta V_n^{(0)}(\Delta u) = o(1)$ . If  $(V_n^{(0)}(\Delta u)) \in U(\mathcal{M})$  is a subsequential convergence condition for  $L$ , then  $(V_n^{(m)}(\Delta u)) \in U(\mathcal{M}^{(m)})$  for each integer  $m \geq 1$  is also a subsequential convergence condition for  $L$ .*

THEOREM 2.2. *For a sequence  $u = (u_n)$ , let  $L - \lim_n u_n = s$  and  $\Delta V_n^{(0)}(\Delta u) = o(1)$ . If  $(V_n^{(0)}(\Delta u)) \in U(\ell_\infty)$  is a subsequential convergence condition for  $L$ , then  $(V_n^{(m)}(\Delta u)) \in U(\ell_\infty^{(m)})$  for each integer  $m \geq 1$  is also a subsequential convergence condition for  $L$ .*

THEOREM 2.3. *For a sequence  $u = (u_n)$ , let  $L - \lim_n u_n = s$  and  $\Delta V_n^{(0)}(\Delta u) = o(1)$ . If  $(u_n) \in U(\mathcal{M})$  is a subsequential convergence condition for  $L$ , then  $(V_n^{(m)}(\Delta u)) \in U(\mathcal{M}^{(m)})$  for each integer  $m \geq 1$  is also a subsequential convergence condition for  $L$ .*

THEOREM 2.4. *For a sequence  $u = (u_n)$ , let  $L - \lim_n u_n = s$  and  $\Delta V_n^{(0)}(\Delta u) = o(1)$ . If  $(u_n) \in U(\ell_\infty)$  is a subsequential convergence condition for  $L$ , then  $(V_n^{(m)}(\Delta u)) \in U(\ell_\infty^{(m)})$  for each integer  $m \geq 1$  is also a subsequential convergence condition for  $L$ .*

To prove these theorems, we need the following lemma and the observation.

LEMMA 2.5 [12]. *Let  $u = (u_n) \in \mathcal{L}$  and  $k, m \geq 0$  be any integers. If  $(V_n^{(k)}(\Delta u)) \in U(\mathcal{A}^{(m)})$ , then  $(n\Delta)_{m+1} V_n^{(k+1)}(\Delta u) = a_n$ , where  $(a_n) \in \mathcal{A}$ .*

*Proof.* If  $(V_n^{(k)}(\Delta u)) \in U(\mathcal{A}^{(m)})$ , it then follows that

$$V_n^{(k)}(\Delta u) = \sigma_n^{(k-1)}(u) - \sigma_n^{(k)}(u) = b_n^{(m)} + \sum_{j=1}^n \frac{b_j^{(m)}}{j} \quad (2.1)$$

for some  $(b_n^{(m)}) \in \mathcal{A}^{(m)}$ . From (2.1), we obtain

$$V_n^{(k-1)}(\Delta u) - V_n^{(k)}(\Delta u) = n\Delta b_n^{(m)} + b_n^{(m)}. \quad (2.2)$$

Subtracting (2.2) from the arithmetic mean of (2.2), we have

$$(V_n^{(k-1)}(\Delta u) - V_n^{(k)}(\Delta u)) - (V_n^{(k)}(\Delta u) - V_n^{(k+1)}(\Delta u)) = b_n^{(m-1)}. \quad (2.3)$$

Equation (2.3) can be expressed as

$$n\Delta V_n^{(k)}(\Delta u) - n\Delta V_n^{(k+1)}(\Delta u) = b_n^{(m-1)}, \quad (2.4)$$

which implies  $(n\Delta)_2 V_n^{(k+1)}(\Delta u) = b_n^{(m-1)}$ . By repeating the same reasoning, we have  $\sigma_n^{(1)}(\omega^{(k+1)}(u)) = (n\Delta)_{m+1} V_n^{(k+1)}(\Delta u) = b_n^{(0)} = b_n$ .  $\square$

For a sequence  $(u_n)$  and for each integer  $m \geq 1$ , we define

$$(n\Delta)_m u_n = n\Delta((n\Delta)_{m-1} u_n), \quad (2.5)$$

where  $(n\Delta)_0 u_n = u_n$  and  $(n\Delta)_1 u_n = n\Delta u_n$ .

*Observation 1* [13]. For each integer  $m \geq 1$ ,

$$\omega_n^{(m)}(u) = (n\Delta)_m V_n^{(m-1)}(\Delta u). \quad (2.6)$$

The proof of Observation 1 easily follows from the mathematical induction.

*Proof of Theorem 2.1.* Assume that  $(V_n^{(0)}(\Delta u)) \in U(\mathcal{M})$  is a subsequential convergence condition for  $L$ . Since  $(V_n^{(0)}(\Delta u)) \in U(\mathcal{M})$ ,  $V_n^{(0)}(\Delta u) = b_n + \sum_{k=1}^n (b_k/k)$  for some  $(b_n) \in \mathcal{M}$ . Hence, we have

$$n\Delta V_n^{(0)}(\Delta u) = n\Delta b_n + b_n. \quad (2.7)$$

Taking the  $(C, 1)$  mean of both sides of (2.7), we obtain  $n\Delta V_n^{(1)}(\Delta u) = V_n^{(0)}(\Delta b) + \sigma_n^{(1)}(b) = b_n$ . Since  $(b_n) \in \mathcal{M}$ ,

$$V_n^{(0)}(\Delta b) = O(1) \quad (2.8)$$

by a result in [5]. Notice that (2.8) can be rewritten as  $V_n^{(0)}(\Delta b) = (n\Delta)_2 V_n^{(2)}(\Delta u) = O(1)$  in terms of the sequence  $u = (u_n)$ . Let  $(V_n^{(m)}(\Delta u)) \in U(\mathcal{M}^{(m)})$ . By Lemma 2.5,  $(\sigma_n^{(1)}(\omega^{(m+1)}(u))) \in \mathcal{M}$ . From the last statement, we conclude that  $\sigma_n^{(1)}(\omega^{(m+2)}(u)) = (n\Delta)_{m+2} V_n^{(m+2)}(\Delta u) = O(1)$ , or equivalently

$$\sigma_n^{(1)}(\omega^{(m+2)}(u)) = (n\Delta)_2 V_n^{(2)}(\Delta \sigma^{(1)}(\omega^{(m-1)}(u))) = O(1). \quad (2.9)$$

It easily follows from the existence of  $L$ -limitability of  $(u_n)$  to  $s$  that

$$L - \lim_n \sigma_n^{(1)}(\omega^{(m-1)}(u)) = 0. \quad (2.10)$$

The condition  $\Delta V_n^{(0)}(\Delta u) = o(1)$  implies that

$$\Delta((n\Delta)_m V_n^{(m)}(\Delta u)) = \Delta V_n^{(0)}(\Delta \sigma^{(1)}(\omega^{(m-1)}(u))) = o(1). \quad (2.11)$$

Taking into account (2.9), (2.10), and (2.11), we obtain that  $(\sigma_n^{(1)}(\omega^{(m-1)}(u)))$  is sub-sequentially convergent. By the fact that every subsequentially convergent sequence is bounded,  $\sigma_n^{(1)}(\omega^{(m-1)}(u)) = O(1)$ , or equivalently

$$\sigma_n^{(1)}(\omega^{(m-1)}(u)) = (n\Delta)_2 V_n^{(2)}(\Delta \sigma^{(1)}(\omega^{(m-4)}(u))) = O(1). \quad (2.12)$$

As in obtaining (2.10) and (2.11), we also have

$$\begin{aligned} L - \lim_n \sigma_n^{(1)}(\omega^{(m-4)}(u)) &= 0, \\ \Delta((n\Delta)_{m-3} V_n^{(m-3)}(\Delta u)) &= \Delta V_n^{(0)}(\Delta \sigma^{(1)}(\omega^{(m-4)}(u))) = o(1), \end{aligned} \quad (2.13)$$

respectively.

Again taking into account (2.12) and (2.13), we obtain that  $(\sigma_n^{(1)}(\omega^{(m-4)}(u)))$  is sub-sequentially convergent. Continuing in this manner, if  $m \equiv 0 \pmod{3}$ , we have that  $((n\Delta)_2 V_n^{(2)}(\Delta u)) = (\sigma_n^{(1)}(\omega^{(2)}(u)))$  is subsequentially convergent and then

$$(n\Delta)_2 V_n^{(2)}(\Delta u) = O(1). \quad (2.14)$$

Since  $L - \lim_n u_n = s$ , we have

$$L - \lim_n \sigma_n^{(1)}(\omega^{(2)}(u)) = 0. \quad (2.15)$$

Again it follows from the conditions  $\Delta V_n^{(0)}(\Delta u) = o(1)$ , (2.14), and (2.15) that  $(u_n)$  is subsequentially convergent.

If  $m \equiv 1 \pmod{3}$ , we have that  $((n\Delta)_0 V_n^{(0)}(\Delta u)) = (V_n^{(0)}(\Delta u)) = (\sigma_n^{(1)}(\omega^{(0)}(u)))$  is sub-sequentially convergent and then

$$V_n^{(0)}(\Delta u) = O(1). \quad (2.16)$$

Clearly, the condition (2.16) implies (2.14).

Again it follows from the conditions  $\Delta V_n^{(0)}(\Delta u) = o(1)$ , (2.14) and (2.15) that  $(u_n)$  is subsequentially convergent.

If  $m \equiv 2 \pmod{3}$ , we conclude that  $(n\Delta V_n^{(1)}(\Delta u)) = (\sigma_n^{(1)}(\omega^{(1)}(u)))$  is subsequentially convergent and then

$$n\Delta V_n^{(1)}(\Delta u) = O(1). \quad (2.17)$$

Clearly, the condition (2.17) implies (2.14).

From the conditions  $\Delta V_n^{(0)}(\Delta u) = o(1)$ , (2.14) and (2.15) it follows that  $(u_n)$  is subsequenceally convergent.  $\square$

*Proof of Theorem 2.2.* Assume that  $(V_n^{(0)}(\Delta u)) \in U(\ell_\infty)$  is a subsequential convergence condition for  $L$ . Since  $(V_n^{(0)}(\Delta u)) \in U(\ell_\infty)$ , by similar calculations in the proof of Theorem 2.1 we have  $(n\Delta V_n^{(1)}(\Delta u)) \in \ell_\infty$ , or equivalently  $n\Delta V_n^{(1)}(\Delta u) = \sigma_n^{(1)}(\omega^{(1)}(u)) = O(1)$ . Let  $(V_n^{(m)}(\Delta u)) \in U(\ell_\infty^{(m)})$ . Then by Lemma 2.5,  $(\sigma_n^{(1)}(\omega^{(m+1)}(u))) \in \ell_\infty$ , or equivalently

$$\sigma_n^{(1)}(\omega^{(m+1)}(u)) = n\Delta V_n^{(1)}(\Delta \sigma^{(1)}(\omega^{(m-1)}(u))) = O(1). \quad (2.18)$$

Since  $L - \lim_n u_n = s$ ,

$$L - \lim_n \sigma_n^{(1)}(\omega^{(m-1)}(u)) = 0. \quad (2.19)$$

The condition  $\Delta V_n^{(0)}(\Delta u) = o(1)$  implies that

$$\Delta((n\Delta)_m V_n^{(m)}(\Delta u)) = \Delta V_n^{(0)}(\Delta \sigma^{(1)}(\omega^{(m-1)}(u))) = o(1). \quad (2.20)$$

Taking into account (2.18), (2.19), and (2.20), we conclude that  $(\sigma_n^{(1)}(\omega^{(m-1)}(u)))$  is subsequenceally convergent, and then  $\sigma_n^{(1)}(\omega^{(m-1)}(u)) = O(1)$ , or equivalently

$$\sigma_n^{(1)}(\omega^{(m-1)}(u)) = n\Delta V_n^{(1)}(\Delta \sigma^{(1)}(\omega^{(m-3)}(u))) = O(1). \quad (2.21)$$

As in obtaining (2.19) and (2.20), we have

$$\begin{aligned} L - \lim_n \sigma_n^{(1)}(\omega^{(m-3)}(u)) &= 0, \\ \Delta((n\Delta)_{m-2} V_n^{(m-2)}(\Delta u)) &= \Delta V_n^{(0)}(\Delta \sigma^{(1)}(\omega^{(m-3)}(u))) = o(1). \end{aligned} \quad (2.22)$$

Taking into account (2.21) and (2.22), we conclude that from the assumption  $(\sigma_n^{(1)}(\omega^{(m-3)}(u)))$  is subsequenceally convergent. Continuing in this manner, if  $m \equiv 0 \pmod{2}$ , we have  $(n\Delta V_n^{(1)}(\Delta u)) = (\sigma_n^{(1)}(\omega^{(1)}(u)))$  is subsequenceally convergent and then,

$$n\Delta V_n^{(1)}(\Delta u) = O(1). \quad (2.23)$$

Since  $L - \lim_n u_n = s$ , we have

$$L - \lim_n \sigma_n^{(1)}(\omega^{(1)}(u)) = 0. \quad (2.24)$$

It follows from the condition  $\Delta V_n^{(0)}(\Delta u) = o(1)$ , (2.23), and (2.24) that  $(u_n)$  is subsequenceally convergent.

If  $m \equiv 1 \pmod{2}$ , we have that  $((n\Delta)_0 V_n^{(0)}(\Delta u)) = (V_n^{(0)}(\Delta u)) = (\sigma_n^{(1)}(\omega^{(0)}(u)))$  is sub-sequentially convergent, and then, we have

$$V_n^{(0)}(\Delta u) = O(1). \quad (2.25)$$

The condition (2.25) implies (2.23).

Taking into account  $\Delta V_n^{(0)}(\Delta u) = o(1)$ , (2.23), and (2.24), we have that  $(u_n)$  is sub-sequentially convergent.  $\square$

*Proof of Theorem 2.3.* Assume that  $(u_n) \in U(\mathcal{M})$  is a subsequential convergence condition for  $L$ . Since  $(u_n) \in U(\mathcal{M})$ , by similar reasoning in the proof of Theorem 2.1, we have  $(V_n^{(0)}(\Delta u)) \in \mathcal{M}$ . Thus, we have  $n\Delta V_n^{(1)}(\Delta u) = O(1)$ . The rest of the proof is as in the proof of Theorem 2.2.  $\square$

*Proof of Theorem 2.4.* Assume that  $(u_n) \in U(\ell_\infty)$  is a subsequential convergence condition for  $L$ . Since  $(u_n) \in U(\ell_\infty)$ , we have  $u_n = b_n + \sum_{k=1}^n (b_k/k)$  for some  $(b_n) \in \ell_\infty$ . Thus  $V_n^{(0)}(\Delta u) = O(1)$ . Let  $(V_n^{(m)}(\Delta u)) \in U(\ell_\infty^{(m)})$ . By Lemma 2.5,  $(\sigma_n^{(1)}(\omega^{(m+1)}(u))) \in \ell_\infty$ , or equivalently

$$\sigma_n^{(1)}(\omega^{(m+1)}(u)) = V_n^{(0)}(\Delta \sigma^{(1)}(\omega^{(m)}(u))) = O(1). \quad (2.26)$$

$L - \lim_n u_n = s$  implies

$$L - \lim_n \sigma_n^{(1)}(\omega^{(m)}(u)) = 0 \quad (2.27)$$

and from  $\Delta V_n^{(0)}(\Delta u) = o(1)$ , we have

$$\Delta((n\Delta)_{m+1} V_n^{(m+1)}(\Delta u)) = \Delta V_n^{(0)}(\Delta \sigma^{(1)}(\omega^{(m)}(u))) = o(1). \quad (2.28)$$

Taking into account (2.26), (2.27), and (2.28), we conclude that  $(\sigma_n^{(1)}(\omega^{(m)}(u)))$  is sub-sequentially convergent, and then  $\sigma_n^{(1)}(\omega^{(m)}(u)) = O(1)$ , or equivalently

$$\sigma_n^{(1)}(\omega^{(m)}(u)) = V_n^{(0)}(\Delta \sigma^{(1)}(\omega^{(m-1)}(u))) = O(1). \quad (2.29)$$

As in obtaining (2.27) and (2.28), we have

$$\begin{aligned} L - \lim_n \sigma_n^{(1)}(\omega^{(m-1)}(u)) &= 0, \\ \Delta((n\Delta)_m V_n^{(m)}(\Delta u)) &= \Delta V_n^{(0)}(\Delta \sigma^{(1)}(\omega^{(m-1)}(u))) = o(1). \end{aligned} \quad (2.30)$$

Again taking into account (2.29) and (2.30), from the assumption we obtain that  $(\sigma_n^{(1)}(\omega^{(m-1)}(u)))$  is sub-sequentially convergent. Continuing in this manner we have that  $(\sigma_n^{(1)}(\omega^{(0)}(u)))$  is sub-sequentially convergent, and then

$$\sigma_n^{(1)}(\omega^{(0)}(u)) = V_n^{(0)}(\Delta u) = O(1). \quad (2.31)$$

Since  $(u_n)$  is  $L$ -limitable to  $s$ , we have

$$L - \lim_n V_n^{(0)}(\Delta u) = 0. \quad (2.32)$$

From the condition  $\Delta V_n^{(0)}(\Delta u) = o(1)$ , (2.31), and (2.32), we conclude that  $(u_n)$  is sub-sequentially convergent.  $\square$

## References

- [1] Č. V. Stanojević, "Analysis of divergence: applications to the Tauberian theory," Graduate Research Seminar, University of Missouri-Rolla, Rolla, Mo, USA, 1999.
- [2] Č. V. Stanojević, *Analysis of Divergence: Control and Management of Divergent Process*, edited by İ. Çanak, Graduate Research Seminar Lecture Notes, University of Missouri-Rolla, Rolla, Mo, USA, 1998.
- [3] G. H. Hardy, *Divergent Series*, The Clarendon Press, Oxford University Press, New York, NY, USA, 1949.
- [4] J. Boos, *Classical and Modern Methods in Summability*, Oxford Mathematical Monographs, Oxford University Press, Oxford, UK, 2000.
- [5] M. Dik, "Tauberian theorems for sequences with moderately oscillatory control modulo," *Mathematica Moravica*, vol. 5, pp. 57–94, 2001.
- [6] F. Dik, "Tauberian theorems for convergence and subsequential convergence with moderately oscillatory behavior," *Mathematica Moravica*, vol. 5, pp. 19–56, 2001.
- [7] A. Tauber, "Ein Satz aus der Theorie der unendlichen Reihen," *Monatshefte für Mathematik und Physik*, vol. 8, no. 1, pp. 273–277, 1897.
- [8] J. E. Littlewood, "The converse of Abel's theorem on power series," *Proceedings of the London Mathematical Society*, vol. 9, no. 2, pp. 434–448, 1911.
- [9] A. Rényi, "On a Tauberian theorem of O. Szász," *Acta Universitatis Szegediensis. Acta Scientiarum Mathematicarum*, vol. 11, pp. 119–123, 1946.
- [10] İ. Çanak and Ü. Totur, "Tauberian theorems for Abel limitability method," submitted for publication.
- [11] İ. Çanak, "Tauberian theorems for a generalized Abelian summability methods," *Mathematica Moravica*, vol. 2, pp. 21–66, 1998.
- [12] İ. Çanak and Ü. Totur, "A note on Tauberian theorems for regularly generated sequences," submitted for publication.
- [13] İ. Çanak and Ü. Totur, "A Tauberian theorem with a generalized one-sided condition," *Abstract and Applied Analysis*, vol. 2007, Article ID 45852, 12 pages, 2007.

İbrahim Çanak: Department of Mathematics, Adnan Menderes University, 09010 Aydın, Turkey  
 Email address: icanak@adu.edu.tr

Ümit Totur: Department of Mathematics, Adnan Menderes University, 09010 Aydın, Turkey  
 Email address: utotur@adu.edu.tr

Mehmet Dik: Department of Mathematics, Rockford College, 5050 E. State Street, Rockford, IL 61108, USA  
 Email address: mdik@rockford.edu